


A Legacy Of Innovation In Hernia Repair



DUALMESH® PLUS

BIOMATERIAL



For use in the reconstruction
of hernias and soft tissue deficiencies
and for the temporary bridging of fascial defects.

The use of prosthetic biomaterials for incisional and ventral hernias larger than four square centimeters greatly reduces the recurrence rate.¹ At the same time, the use of a biomaterial can increase the infection rate. Postoperative infection has been shown to be a significant factor in hernia recurrence. Studies have shown that it takes fewer organisms to produce an infection if biomaterials are present.

Traditionally, surgeons have used various prophylactic regimens when a prosthetic material is used. These have had limited results in reducing the incidence of infection. This may be especially true in the repair of recurrent hernias where a significant increase in the incidence of infection has been demonstrated. Additional individual patient characteristics (i.e., diabetes, ascites, steroid therapy) may also increase the risk of infection.

W. L. Gore & Associates, Inc — the innovator and leader in expanded polytetrafluoroethylene (ePTFE) technology — has combined an innovative surgical biomaterial with two antimicrobial preservative agents. GORE DUALMESH® PLUS Biomaterial features not only the anti-microbial agents of chlorhexidine diacetate and silver carbonate, but also an advanced tissue ingrowth surface.

The two antimicrobial preservatives act synergistically to inhibit colonization of the device and resist initial biofilm formation on the device for up to 14 days post implantation. Zone-of-inhibition bioassays have found that this device has substantial preservative activity against the following gram-positive and gram-negative organisms:

- ▶ Methicillin-resistant Staphylo-coccus aureus (MRSA)
- ▶ Vancomycin-resistant Entero-coccus faecalis (VRE)
- ▶ Staphylococcus epidermidis
- ▶ Klebsiella pneumoniae
- ▶ Pseudomonas aeruginosa
- ▶ Candida albicans
- ▶ Escherichia coli
- ▶ Staphylococcus aureus
- ▶ Group A Streptococcus
- ▶ Acinetobacter baumannii

Gore's antimicrobial technology is coupled with the CORDUROY® Tissue Ingrowth Surface of the GORE DUALMESH® PLUS Biomaterial. In addition to a low-porosity, smooth visceral surface, the fascial interface side features ePTFE "ridges" and "valleys." Experimental models have shown that this surface stimulates a heightened tissue fixation process that results in a firm bond to host tissue². Additionally, the processing of the GORE DUALMESH® PLUS Biomaterial with the anti-microbial agents results in a darkening of the barrier membrane side to provide for side differentiation and to reduce glare from laparoscopic light sources.

GORE CORDUROY® Surface for tissue ingrowth





DUALMESH® PLUS

BIOMATERIAL

Surface Orientation

Proper surface orientation is essential for the GORE DUALMESH® PLUS Biomaterial to function as intended. The smooth, non-ingrowth surface is shaded darker in comparison to the textured ingrowth surface to provide side differentiation and to reduce glare from laparoscopic light sources. The darker shaded surface should be placed adjacent to tissues where minimal tissue attachment is desired. The lighter shaded surface has an open microstructure that stimulates tissue ingrowth and should be placed adjacent to tissues where incorporation is desired.

Suture/Staple Recommendations

- Use only nonabsorbable sutures, such as GORE-TEX® Suture, with a noncutting needle (such as taper or piercing point). For best results, use monofilament sutures.
- Suture size should be determined by surgeon preference and the nature of the reconstruction. A bite and spacing ratio of 1:1 is recommended.³
- Staples or helical tacks (also known as helical coils) can be used as an alternative to sutures. Staple size and staple or tack spacing should be determined by surgeon preference.

Surgical Drains/Seroma

- Use of a drain should reflect surgeon preference.^{4,5} Closed-suction drains rather than gravity drains are recommended to prevent handling-related infections.
- In any hernia defect repair it is possible for seroma to occur up to six weeks postoperatively. Aspiration or placement of a drain, followed by pressure dressing, may resolve the seroma.^{6,7,8,9}


Use in a Contaminated Field/Postoperative Infection

- GORE DUALMESH® PLUS Biomaterial is not recommended for use in grossly infected tissue.
- Appropriate preoperative and postoperative use of local and systemic antibiotics is highly recommended. In the event of a postoperative infection, an aggressive regimen of antibiotic treatment, possibly including antibiotic irrigation, aspiration and debridement of the affected area may resolve the infection. Persistent infection may necessitate removal of the device.


Open Healing

- When using this device as a temporary external bridging device where primary closure is not possible, use measures to avoid contamination. The entire device should be removed as early as clinically feasible, not to exceed 45 days after placement.
- When using this device as a permanent implant and unintentional exposure occurs, treat to avoid contamination, or device removal may be necessary.

CONTRAINDICATIONS: Patients with hypersensitivity to chlorhexidine or silver; reconstruction of cardiovascular defects; reconstruction of central nervous system or peripheral nervous system defects; pre-term and neonatal populations. **WARNINGS:** Use with caution in patients with methemoglobinopathy or related disorders. When used as a temporary external bridging device, use measures to avoid contamination; the entire device should be removed as early as clinically feasible, not to exceed 45 days after placement. When unintentional exposure occurs, treat to avoid contamination or device removal may be necessary. Improper positioning of the smooth non-textured surface adjacent to fascial or subcutaneous tissue will result in minimal tissue attachment. **POSSIBLE ADVERSE REACTIONS:** Contamination, infection, inflammation, adhesion, fistula formation, seroma formation, hematoma and recurrence.



GORE DUALMESH® PLUS Biomaterial inoculated with *Staphylococcus aureus* in a rabbit model 10 days post-inoculation. H&E stain showing large numbers of acute inflammatory cells and eosinophilic cellular debris (arrows) covering the surface of the implant. The interstices of the implant (I) contain small amounts of eosinophilic cellular debris and very few acute inflammatory cells suggesting protection of the implant interstices from bacterial infection. H&E; 20x magnification



GORE DUALMESH® Biomaterial inoculated with *Staphylococcus aureus* in a rabbit model 10 days post-inoculation. H&E stain showing large numbers of acute inflammatory cells and necrotic cellular debris covering the surface of the implant (arrows). The interstices of the implant (I) also contain large numbers of acute inflammatory cells and eosinophilic cellular debris indicating bacterial infection of the implant surface and interstices. H&E; 20x magnification

Sizes Available

CATALOGUE NUMBER		NOMINAL WIDTH X LENGTH
(1 mm)	(2 mm)	
1DLMCP02	–	8 cm x 12 cm
1DLMCP03	1DLMCP200	10 cm x 15 cm*
1DLMCP04	1DLMCP201	15 cm x 19 cm*
1DLMCP05	–	7.5 cm x 10 cm
1DLMCP06	1DLMCP202	18 cm x 24 cm
1DLMCP07	1DLMCP203	20 cm x 30 cm
1DLMCP08	1DLMCP204	26 cm x 34 cm*
1DLMCP11	–	10 cm x 30 cm**

* oval shaped

** trapezoid shaped

Packaged Sterile

References

- Hesselink VJ, Luijendijk RW, de Wilt JHW, Heide R, Jeekel J. An evaluation of risk factors in incisional hernia recurrence. *Surgery, Gynecology & Obstetrics* 1993;176:228-234.
- LeBlanc KA, Bellanger D, Rhynes KV, Baker DG, Stout RW. Tissue attachment strength of prosthetic meshes used in ventral and incisional hernia repair. A study in the New Zealand white rabbit adhesion model. *Surgical Endoscopy* 2002;16(11):1542-1546.
- Nealon TF. *Fundamental skills in surgery*. Philadelphia: Saunders, 1979:47.
- Nyhus LM, Condon RE, eds. *Hernia*. 4th ed. Philadelphia: Lippincott, 1995:331-6.
- Hamer-Hodges DW, Scott NB. Replacement of an abdominal wall defect using expanded PTFE sheet (GORE-TEX). *J R Coll Surg Edinb* 1985;30:65-7.
- Ponka JL. *Hernias of the abdominal wall*. Philadelphia: Saunders, 1980:339, 352, 392.
- Durden JG, Pemberton LB. Dacron mesh in ventral and inguinal hernias. *Am Surg* 1974;40:662-5.
- Reisfeld D, Schechner R, Wetzel W. Traumatic lumbar hernia. *Surg Rounds* 1989 Mar;12:69-72.
- Nichter LS, Morgan RF, Dufresne CR, Lambruschi P, Edgerton MT. Rapid management of persistent seromas by sclerotherapy. *Ann Plast Surg* 1983;11:233-6.

W. L. Gore & Associates, Inc.
Flagstaff, AZ 86004

800.437.8181 (US)
928.779.2771 (US)
00800.6334.4673 (EU)

For international contacts and additional product information, visit goremmedical.com